

# Documentos de trabajo PROESA



**Young Latino Children Obesity: The Importance of Socioeconomic Status,  
Home Language, TV watching, and Physical Activity**

Claudia L. Galindo, PhD & Sergio I. Prada, MPA, PhD  
Documentos PROESA #12

**Young Latino Children Obesity: The Importance of  
Socioeconomic Status, Home Language, TV Watching, and  
Physical Activity**

**Claudia L. Galindo, PhD & Sergio I. Prada, MPA, PhD**

**Documentos PROESA #12**

## DOCUMENTOS PROESA

ISSN: 2256-4128

Enero de 2017

Comité Editorial:

Ramiro Guerrero, Ms

Sergio I. Prada, PhD

Victoria E. Soto, PhD

Yuri Takeuchi, MD

PROESA – Centro de Estudios en Protección Social y Economía de la Salud  
Cl 18 # 122-135 Universidad Icesi Casa Rocha segundo piso  
Tel: +57(2)3212092 Cali Colombia  
[www.icesi.edu.co/proesa](http://www.icesi.edu.co/proesa)  
[www.proesa.org.co](http://www.proesa.org.co)

# Young Latino Children Obesity: The Importance of Socioeconomic Status, Home Language, TV Watching, and Physical Activity

Claudia L. Galindo, PhD<sup>1</sup> & Sergio I. Prada, MPA, PhD<sup>2</sup>

## Abstract

The incidence of young Latino children obesity is a public health concern. Among 5-year-olds, 16% of Latinos are obese compared with 10% of whites. Obesity disparities between non-Latino white and Latino children increase significantly with age. The purpose of this study is to examine the importance of socioeconomic status (SES), home language, physical activity, and TV watching for explaining obesity disparities between 5-year-old white and Latino children. Also, the long-lasting influence of these four variables is studied by analyzing their impact on obesity as children grew to ages 6, 8, and 10. We used a nationally representative sample drawn from the Early Childhood Longitudinal Study-Kindergarten Cohort (ECLS-K) collected in 1998-99. Logistic regression models with adjusted standard errors for the complex sampling design and nested structure of the ECLS-K were estimated. Analyses were conducted in 2012. SES and home language were significantly associated with obesity after controlling for covariates. Similarly, physical activity had a negative association and TV watching has a positive association with obesity. When accounting for all four variables, obesity disparities between white and Latino children became statistically non-significant. All four factors had lasting effects on obesity even as children grew to ages 6, 8, and 10. Findings identified mechanisms to reduce young Latino children' obesity disparities. The benefits of these mechanisms persist throughout early adolescence, which is particularly important given the need to implement interventions with long term effects.

**Keywords:** Children obesity, Latinos, socioeconomic status, home language, physical activity.

---

<sup>1</sup> Associate Professor, College of Education, University of Maryland, College Park.

<sup>2</sup> Director at PROESA & Professor Universidad ICESI, Facultad Ciencias Administrativas y Económicas, Departamento de Economía.

# Obesidad en niños latinos: la importancia del estatus socioeconómico, el lenguaje en casa, consumo de televisión, y la actividad física.

Claudia L. Galindo, PhD<sup>3</sup> & Sergio I. Prada, MPA, PhD<sup>4</sup>

## Resumen

La incidencia en la obesidad en niños jóvenes latinos es un problema de salud pública. Entre los niños de 5 años de edad, el 16% de los latinos son obesos en comparación con 10% de los blancos. Las brechas en la obesidad entre los niños no latinos y los blancos se incrementa significativamente con la edad. El propósito de este estudio es examinar la importancia del estatus socioeconómico (ESE), la lengua materna, la actividad física, y el consumo de televisión como explicación de las brechas de obesidad entre niños blancos de 5 años de edad y niños latinos. A su vez, la influencia de estas cuatro variables en el mediano plazo es estudiada analizando su impacto en la obesidad cuando los niños crecen a edades de los 6, 8 y 10 años. Usamos la muestra nacional representativa Early Childhood Longitudinal Study-Kindergarten Cohort (ECLS-K) tomada en 1998-99 en los Estados Unidos. Modelos de regresiones logísticas con errores estándares ajustados por el diseño complejo de muestreo y la estructura anidada de ECLS-K fueron estimados. Los análisis fueron hechos en 2012. Los resultados muestran que ESE y la lengua maternal están asociados significativamente con la obesidad tras controlar por otras variables. Similarmente, la actividad física tiene una asociación negativa y el ver televisión tiene una asociación positiva con la obesidad. Cuando incluimos las cuatro variables, la brecha en obesidad entre niños blancos y latinos se vuelve estadísticamente no significativa. Todos los cuatro factores tienen efectos duraderos en la obesidad cuando se observan a las edades de 6, 8 y 10 años. Los hallazgos identifican mecanismos para reducir la brecha de los niños jóvenes latinos. Los beneficios de esos mecanismos persisten a través de la temprana adolescencia, la cual es particularmente importante dada la necesidad de implementar intervenciones de largo plazo.

**Keywords:** Obesidad infantil, Latinos, estatus socioeconómico, lenguaje en casa, actividad física.

---

<sup>3</sup> Associate Professor, College of Education, University of Maryland, College Park.

<sup>4</sup> Director de PROESA & Profesor Universidad ICESI, Facultad Ciencias Administrativas y Económicas, Departamento de Economía.

## CONTENIDO

<b>1 INTRODUCTION</b> .....	<b>6</b>
1.1 EXPLAINING THE OBESITY EPIDEMIC AMONG LATINO CHILDREN .....	6
<b>2. METHODS</b> .....	<b>8</b>
2.1 STATISTICAL METHODS.....	9
<b>3. RESULTS</b> .....	<b>10</b>
3.1 ASSOCIATION BETWEEN SES, HOME LANGUAGE, AND OBESITY .....	10
3.2 THE IMPORTANCE OF TV WATCHING AND PHYSICAL ACTIVITY .....	10
3.3 LONG-LASTING EFFECTS OF SES, HOME LANGUAGE, TV WATCHING AND PHYSICAL ACTIVITY .....	11
3.4 LIMITATIONS .....	11
<b>4. DISCUSSION AND CONCLUSIONS</b> .....	<b>11</b>
<b>5. REFERENCES</b> .....	<b>14</b>

## 1 Introduction

The progression of the obesity epidemic across different age groups in the United States has been well chronicled. In the past 30 years, obesity rates doubled from 5% to 10.4% for children ages 2-5, tripled from 6.5% to 19.6% for ages 6-11, and almost quadrupled from 5% to 18.1% for ages 12-19.<sup>1</sup> The incidence of obesity is even more alarming for racial/ethnic minority children, with blacks, Latina/os, and Native Americans showing the highest rates of weight problems.<sup>2-7</sup> Minority children's weight disadvantages are observed as early as the age of 5. At the start of kindergarten, about 16% of Latino children are obese, compared with only 10% of white children.<sup>8</sup> Furthermore, obesity disparities increase disproportionately for minority children with age and persist in adolescence<sup>9-12</sup> and adulthood.<sup>13</sup> Thus, although obesity is a significant health problem across all children, it is more concentrated among racial/ethnic minority children, who are more likely to experience socioeconomic disadvantages.<sup>14</sup>

### 1.1 Explaining the Obesity Epidemic among Latino children

Sociologists apply two main theoretical perspectives to explain health disparities and the incidence of obesity among Latino children: cultural and structural/economic. By combining these two theoretical approaches a holistic understanding of a major public health concern is achieved. Cultural arguments examine how group-specific cultural beliefs and family practices may impact weight problems. From this perspective, the sociocultural influences on childhood obesity, including parents' beliefs about well-being, body perceptions, and the meaning and function of food and eating, are key factors in understanding the obesity problem of Latino children.<sup>15</sup> Research, for example, shows that perceptions of body size and acknowledgment of weight problems are different across ethnic/racial groups, with minorities considering a larger body size as ideal and healthy than do whites.<sup>15-17</sup> For some racial/ethnic minority females, having heavy infants implies having healthy babies<sup>17</sup> and being heavy is considered a sign of strength, high self-esteem, and attractiveness.<sup>18,19</sup>

Furthermore, some food habits and customs are at the center of racial/ethnic minorities' value system and are major part of many Latino parents' cultural identity.<sup>20,21</sup> In a study with Latina mothers, Sussner and her colleagues (2008) found that mothers revealed the cultural importance of "finishing their plates."<sup>21</sup> Also, cultural differences were associated with perceptions of healthiness of different types of food.<sup>22</sup> Cultural influences not only impact the way adults eat, but also how they socialize children around food behaviors and use food as a tool for parenting to reward behaviors.<sup>17</sup>

Immersed within discussions of cultural explanations of minority students' obesity are questions related to the role that acculturation – defined as the process of becoming a member of the U.S. culture and society through the adoption of practices and styles of the dominant group<sup>23</sup> – plays on childhood obesity.<sup>22</sup> This is particularly important for Latinos, about half (54%) of whom are foreign-born and a significant proportion are U.S.-born but live in immigrant families.<sup>24</sup> Research has shown inconsistent findings regarding the role acculturation

plays on Latinos' health outcomes. On the one hand, research shows that less acculturated Latino families have healthier eating habits<sup>21, 25</sup> and healthier prenatal practices.<sup>26</sup> And, as the length of residence in the United States increases, so do smoking and weight gains for Latino adolescents.<sup>27</sup> These findings support the "Immigrant Paradox," which argues that although immigrant families face financial challenges, foreign-born or first generation children often have better outcomes (e.g., health and education) than their native-born counterparts. Yet, additional research shows improvement of outcomes over time or in subsequent generations suggesting worst results for the first generation.<sup>28, 29</sup>

Sociological structural theories propose a different perspective on health disparities and childhood obesity. These theories (e.g. structural-functionalism approach<sup>30</sup>, social Marxism<sup>31</sup>) argue that individual outcomes are being shaped by an underlying social and economic structure<sup>32</sup> and therefore the position of racial/ethnic minority groups within the U.S. social hierarchy and their sustained economic disadvantages are the main determinants of their health problems. Structural accounts of minorities' obesity focus on the impact of poverty and low income through which poverty perpetuates health disparities. On average, racial/ethnic minorities, especially African Americans and Latinos, have lower socioeconomic status (SES) and fewer material resources than whites. In 2010, poverty rates for Latinos and blacks were around 27%, compared to 9% for non-Latino whites. Family median incomes for Latinos and blacks were \$37,000 and \$32,000, respectively, which represents only around half the median income of whites.<sup>33</sup>

The importance of poverty in accounting for weight problems is vastly recognized in the literature since children and youth living in poverty are more likely to be obese and experience other health problems than those with more economic resources.<sup>10, 34</sup> The negative impact of childhood poverty on weight has long-lasting effects as one study (Wells, Evans, Beavis, and Ong 2010) implemented in upstate New York found that poverty negatively affected weight even when children reached the ages of 13 to 17.<sup>35</sup> Similar findings are reported by Poulton and his colleagues (2002), who found childhood SES negatively associated with obesity as well as with dental and mental health issues up to the age of 26 years old.<sup>36</sup>

Research consistently shows that, on average, children living in poverty have less access to recreational activities and time for exercising,<sup>37</sup> are less likely to exercise and be physically active,<sup>4</sup> and are more likely to watch TV.<sup>17</sup> Levels of poverty are also related to increased access to fast food restaurants,<sup>38</sup> food insecurity,<sup>39</sup> and lack of access to healthy foods.<sup>40</sup> Yet the role of SES in explaining young Latino children's obesity is contested. Balistri and Van Hook (2009), using the Early Childhood Longitudinal Study-Kindergarten Cohort (ECLS-K), found a negative association between parents' education and BMI only for Latino children with foreign-born parents.<sup>41</sup> They found no association between these variables for Latino children with U.S.-born parents. Similar findings were obtained in a study with only Mexican-origin adolescents.<sup>42</sup> The inconsistent evidence on the relationship between SES indicators and obesity may be related to interaction effects between these variables and other variables like

gender,<sup>43</sup> or related to the fact that higher SES does not seem to protect Hispanic children against obesity.<sup>44</sup>

The purpose of this study was to examine the influence of home language and SES, as proxy of structural and acculturation indicators, on obesity disparities between non-Latino white and Latino young children. A commonly used indicator of acculturation in health and psychology research is language patterns.<sup>45</sup> Then, the joint influence of TV watching and physical activity on obesity is examined. It is also important to disentangle whether these micro-level mechanisms explain the effects of home language and SES on the same dependent variable. We know that TV watching and physical activity are separately associated with weight problems.<sup>46-48</sup> We also know that TV watching is associated with exposure to advertisements of high calorie foods, increased consumption of high-calorie foods and lower levels of physical activity.<sup>8</sup> Understanding the role of TV watching and physical activities on obesity disparities may be particularly important since Latino children are more likely to watch TV and are less likely to be physically active than white children.<sup>49,50</sup> At the same time, Latino parents report positive attitudes to TV watching and experience significant barriers to supporting their children in being physically active.<sup>48,51</sup> We analyzed the impact of these variables over time, when children were 5-years-old, on average, and then when they turned 6, 8 and 10 years old.

## 2. Methods

The data used in this study came from the Early Childhood Longitudinal Study (ECLS-K) Kindergarten Class of 1998-99, sponsored by the National Center of Education Statistics. The ECLS-K focused on young children's well-being outcomes and collected information through parent, teacher, and administrator interviews between the start of kindergarten through eighth grade. Using a multistage probability sampling design, the ECLS-K includes a nationally representative sample of about 21,000 kindergarteners in over 1,000 schools (for methodological details, see NCES, 2002).<sup>52</sup> In this article, the information analyzed came from Fall 1998 and Spring 1999 waves of data collection. To analyze long-term effects, follow-up obesity measures from Spring 2000, 2002, and 2004 were also included in the analysis. Analyses were conducted in 2012.

The study sample was restricted only to native non-Latino whites and Latino children (roughly 11,470 children). The sample was further restricted to only children deemed as obese or with normal weight based on CDC's tables of BMI-for-age growth charts percentile rankings (roughly 9,000 children). The sample for this study included 70% native non-Latino whites and 30% Latinos of any race. Almost half of the sample (48%) were females, 22% lived in single-parent families and the average number of siblings at home was 1.46. About 78% of the children lived in only English-speaking homes and 36% of their parents had a high school degree or lower education. The sample also included 14% obese and 86% normal-weight children.

The primary dichotomous outcome measure is obesity. To classify children as obese ( $y=1$ ) or with normal weight ( $y=0$ ) the body mass index (BMI) and the CDC's tables of BMI-for-age growth charts percentile rankings were used. Children were deemed as obese if their percentile rankings were equal to or greater than the 95<sup>th</sup> percentile. Children were considered as with normal weight if their percentile rankings were higher or equal to 5<sup>th</sup> percentile through 85<sup>th</sup> percentile.<sup>53</sup> BMI was calculated based on weight and height collected by ECLS-K staff using a Shorr board and a digital scale. Obesity was measured at four points in time: when children were on average 5-years-old (spring of kindergarten), and when children aged to 6 (spring of first grade), 8 (spring of third grade), and 10 (spring of fifth grade).

The main explanatory variables of the study were SES, home language, physical activity, and TV watching. To measure SES, a composite scale (mean=1 and a standard deviation=0) was used. The scale was included in the ECLS-K data and was created based on child's mother's and father's educational attainment and occupation, and family income.<sup>52</sup> To measure home language, parents' reported the language(s) that the mother and father spoke to the student (and vice versa). Children were defined as living in homes where language used was reported as 1) only English (reference category), 2) English and Spanish, or 4) only Spanish. TV watching was measured from parents' reported number of hours the child spent watching either TV or videos on school days. Physical activity was measured using an ordinal variable indicating amount of aerobic exercise that the child did on a regular basis. Parents reported whether their child exercised 1) less than other children, 2) about the same or 3) more than other children.

## 2.1 Statistical Methods

Logistic regression models with stepwise variable selection were estimated with adjusted standard errors given the complex sampling design and the nested structure of the ECLS-K using *svy* commands in Stata-12. Results were reported in odds-ratio. Four sequential regression models (Table 1) were estimated to analyze the relationship between obesity (Spring 1999) as the dependent variable and key explanatory variables for young children. *Model 1* included race/ethnicity (Latino=1 and white=0) as the independent variable. *Model 2* added SES and home language as independent variables. *Model 3* added physical activity and TV watching and excluded home language and SES. *Model 4* includes all four independent variables. To analyze the long-lasting impact of these variables (Table 2), three models similar to *Model 4* were estimated, but with obesity measured when children turned 6 (*Model 5*), 8 (*Model 6*), and 10 years old (*Model 7*) as the dependent variables. The following specific confounders were included as control variables in all models: weight at birth in pounds, gender, number of siblings at home, single-parent family, mother's age in years and depression.

### 3. Results

#### 3.1 Association between SES, Home Language, and Obesity

As previous research has shown, young Latino children had higher incidence of obesity than white children (Model 1, Table 1). Latinos had two times the odds as whites to be obese, after controlling for covariates (weight at birth, gender, siblings at home, single-parent family, and mother's age and depression). The association between obesity and SES and home language was presented in Model 2, Table 1. After controlling for covariates, both variables were significantly associated with obesity. For a one-unit increase in SES, the expected change in log odds was 0.65. Similarly, compared to children living in only-English speaking homes, children living in homes with different language environments had higher incidences of obesity. Children living in only-Spanish speaking homes had 1.48 times the odds than those in only-English speaking homes to be obese.

When the influence of SES and home language was taken into account, the obesity gap between white and Latino children decreased considerably, although it remained borderline statistically significant ( $p$ -value=0.05).

#### 3.2 The importance of TV watching and physical activity

We then examined the impact of hours of TV watching and amount of aerobic exercise (Models 3 and 4 in Table 1). Children whose parents reported they participated more in aerobic exercises than their peers and watched less TV were less likely to be obese. The significance of these findings were independent of including or not home language and SES in the model. After controlling for covariates and home language and SES, for a one-hour increase in TV watching, the expected change in log odds was 1.09. Also, the difference in log of the odds between a child who exercises more than other children in comparison to a child with lower levels of exercise is 0.67.

When we took into account the influence of TV watching and aerobic physical activity, together with home language and SES, the obesity gap between white and Latino young children became statistically non-significant ( $p$ -value=0.08). The difference in obesity between the two racial/ethnic groups was explained by including all four variables in the model. These findings are specifically relevant to reduce the incidence of obesity among young Latino children because these children watch more TV and participate less in aerobic exercise than white children. On average, young Latino children in this sample watched two hours of TV a day whereas white children watched only 1.7 hours ( $t$ -test= 13.44,  $p$ -value <0.00). Also, a lower proportion of Latino children (18%) than of white children (21%) are participating more than other children in aerobic exercise ( $\chi^2$ =35.58,  $p$ -value<0.00). Nevertheless, it is important to mention that physical activity and TV watching did not explain the SES or home language effects suggesting that other factors need to be explored.

### **3.3 Long-lasting effects of SES, Home Language, TV watching and physical activity**

In order to develop interventions to reduce the young Latino obesity epidemic, it is important to identify mechanisms that have long-lasting effects and whose benefits are sustained over time. As Table 2 shows, the positive association between obesity and participation in aerobic exercises and the negative association between obesity and number of hours watching TV remained statistically significant even when children grew to ages 6, 8 and 10. Long-lasting effects of SES and home language were also observed over time. The associations observed across time maintain similar levels of significance and magnitude of associations.

### **3.4 Limitations**

Future studies need to use improved acculturation measures. XX and colleagues argued that bidimensional (mutual influence of both cultures) and multidimensional better capture the complexity and dynamism of this construct. Home language, although a commonly used proxy, may not be able to capture this complexity. This limitation should be considered when interpreting the findings of this paper. Additional studies are also needed to examine the specific mechanisms that neutralize Latino young children's obesity disadvantages, as well as the effectiveness of interventions to reduce health problems. These future studies should include stronger measures of physical activity and exercise. The measures used in this study were based on parents' reports, which could be problematic given that it is a subjective measurement easily influenced by parents' perceptions. In the same way, we need research that takes into account the diversity of the Latino population in terms of country of origin<sup>8</sup> and gender<sup>43</sup> given that incidence of obesity varies function of subgroup memberships.

Additional limitations of this study include the fact that a secondary data was used and therefore important constructs (e.g., parents' obesity, time-varying exercise and TV watching patterns) that are also associated with the outcomes of interest. Also, these findings apply only to obese and normal children. Future studies should focus on examining patterns and influences for children who are at risk of becoming obese (overweight children).

## **4. Discussion and Conclusions**

This study contributes new knowledge on Latino young children's health inequalities by analyzing the importance of SES, home language, physical activity, and TV watching for explaining obesity disparities between 5-year-old white and Latino children using nationally representative data, the ECLS-K. This paper also studied the lasting effect of these variables as children grow to age 6, 8, and 10. Given the high incidence of obesity among Latino children and its negative consequences on different dimensions of well-being, it is imperative to identify malleable mechanisms that could contribute to the reduction of health disparities among these children.

The present study provides empirical support for the importance of considering both structural/economic and cultural explanations when examining health disadvantages for minority groups. As Patterson (2000) argued, structural factors condition the context in which cultural attributes are manifested therefore both frameworks need to be examined interactively.<sup>54</sup>

Results showed that socioeconomic status and home language were relevant factors in explaining Latino children's obesity disparities after taking into account confounders. As other research has shown, children with greater access to economic resources were less likely to be obese than those that had fewer access to resources.<sup>10</sup> Also, children living in English-speaking homes were less likely to be obese than children living in homes with other language environments. Home language is considered as one indicator of acculturation, perhaps the most important one,<sup>55</sup> given that the language that children and their families use at home may reflect a tendency to maintain group-specific cultural values, norms, and practices. This research suggests that it may be possible that some practices and customs that Latino families have had in the past may not be conducive to healthy weight in the U.S. context. This is consistent with research conducted with Latino adolescents.<sup>56</sup> In both cases, acculturation does not seem to be as much of a risk factor as other research on health outcomes among immigrants have shown.<sup>57</sup> Future studies are needed to better understand the mechanisms through which and the context in which acculturation impacts obesity among Latino young children. Understanding the relationship between acculturation and obesity is particularly important for Latinos given their high proportion of foreign-born and those living in immigrant families.<sup>24</sup>

SES and home language are statistically significant predictors of obesity, yet these variables are not easily malleable and therefore are not particularly relevant to inform the development of interventions to reduce children's obesity. The results of this study show that increasing Latino children physical activity and reducing hours watching TV have important positive consequences for reducing Latino and white children disparities in obesity. The benefits of increasing physical activity and reducing TV watching have long-lasting effects even when children turn 6, 8, and 10 years old, which is particularly important given the need to implement interventions that will sustain their benefits over time.

Challenges remain to developing interventions to reduce TV watching and to increase physical activity for Latino young children. Latino children are more likely to live in poor neighborhoods and face lack of safety and higher violent crime than white children.<sup>58</sup> These are important barriers to outdoor physical activity for adults<sup>59</sup> and for adolescent girls.<sup>60</sup> Perception of neighborhood insecurity may be even more important for young children's physical activity. Also, TV may have additional roles for Latino families besides entertaining, as it was shown in a study with Latino mothers who emphasize the importance of TV as a tool to learn English for them and their children.<sup>51</sup>

This study provides a useful theoretical framework to understand young Latino obesity by integrating the explanatory power of both, cultural and structural/economic frameworks. Findings from this study also support the relevance of macro-contextual factors captured through SES and home language proxies. Furthermore, increasing physical activity and reducing TV watching, as moldable micro-level mechanisms, have important implications for reducing health inequalities and developing interventions for Latino young children, independently of the influence of acculturation and SES factors.

### **Acknowledgements**

This study was funded by the Robert Wood Johnson Foundation through its national program, Salud America! (#66465). The RWJF Research Network to Prevent Obesity Among Latino Children ([www.salud-america.org](http://www.salud-america.org)) Salud America!, led by the Institute for Health Promotion Research at The University of Texas Health Science Center at San Antonio, Texas, unites Latino researchers and advocates seeking environmental and policy solutions to the epidemic. We appreciate the excellent research assistance of Amy Pucino.

No financial disclosures were reported by the authors of this paper.

## 5. References

1. Ogden C, Carroll M. Prevalence of obesity among children and adolescents: United States, trends 1963-1965 through 2007-2008. National Center for Health Statistics (NCHS) Health E-Stat, 2010. Available from: [http://www.cdc.gov/nchs/data/hestat/obesity\\_child\\_07\\_08/obesity\\_child\\_07\\_08.pdf](http://www.cdc.gov/nchs/data/hestat/obesity_child_07_08/obesity_child_07_08.pdf).
2. Detar A, Strum R, Magnabosco J. Childhood overweight and academic performance: National study of kindergartners and first-graders. *Obes Res* 2004; 12(1): 58-68.
3. Kimbro R, Brooks-Gunn J, McLenahan S. Racial and ethnic differentials in overweight and obesity among 3-year old children. *Am J Public Health* 2007;97(2): 298-305.
4. Rose D, Bodor N. Household food insecurity and overweight status in young school children: Results from the early childhood longitudinal study. *Pediatr* 2006;117(2): 464-73.
5. Maher EJ, Li G, Carter L, Johnson DB. Preschool child care participation and obesity at the start of kindergarten. *Pediatr* 2008;122(2): 322-30.
6. Singh GK, Kogan MD, Van Dyck PC, Siahpush M. Racial/ethnic, socioeconomic, and behavioral determinants of childhood and adolescent obesity in the United States: Analyzing independent and joint associations. *Ann Epidemiol* 2008;18(9): 682-95.
7. Ogden CL, Carroll MD, Curtin LR, Lamb MM, Flegal KM. Prevalence of high Body Mass Index in US children and adolescents, 2007–2008. *J Am Med Assoc* 2010; 303(3) :242–9.
8. Galindo C, Prada S, Shinogle J, Kirk A. Obesity among young Latino children: Disparities and changes over time. *SaludAmerica! The Robert Wood Johnson Foundation Research Network to Prevent Obesity Among Latino Children*, 2011. Available from: <http://www.umbc.edu/mipar/Documents/Obesity.pdf>.
9. Giammattei J, Blix G, Marshak HH, Wollitzer AO, Pettitt DJ. Television watching and soft drink consumption: Associations with obesity in 11- to 13-year-old-schoolchildren. *Arch Pediatr Adolesc Med* 2003;157(9): 882-86.
10. Babey S, Hastert T, Wolstein J, Diamant A. Income disparities in obesity trends among California adolescents. *Am J Public Health* 2010;100(11): 2149-55.

11. Richmond TK, Subramanian SV. School level contextual factors are associated with the weight status of adolescent males and females. *Obesity* 2008;16(6): 1324-30.
12. Schaefer SE, Salazar M, Bruhn C, Saviano D, Boushey C, Van Loan MD. Influence of race, acculturation, and socioeconomic status on tendency toward overweight in Asian-American early adolescent females. *J Immigr Minor Health* 2009; 11(3): 188-197.
13. Chang VW, Hillier AE, Mehta NK. Neighborhood racial isolation, disorder and obesity. *Soc Forces* 2009;87(4): 2063-92.
14. DeNavas-Walt C, Proctor B, Smith J. Income, poverty, and health insurance coverage in the United States: 2010. United States Census Bureau, 2011. Available from: <http://www.census.gov/prod/2011pubs/p60-239.pdf>.
15. Kaufman L, Karpati A. Understanding the sociocultural roots of childhood obesity: Food practices among Latino families of Bushwick, Brooklyn. *Soc Sci Med* 2007;64(11): 2177-88.
16. Fitzgibbon ML, Blackman LR, Avellone ME. The relationship between body image discrepancy and Body Mass Index across ethnic groups. *Obes Res* 2000;8(8): 582-9.
17. Kumanyika SK. Environmental influences on childhood obesity: Ethnic and cultural influences in context. *Physiol Behav* 2008;94(1): 61-8.
18. Davidson M, Knafl KA. Dimensional analysis of the concept of obesity. *J Adv Nurs* 2006;54(3): 342-50.
19. Contento IR, Basch C, Zybert P. Body image, weight, and food choices of Latina women and their young children. *J Nutr Educ Behav* 2003;35(5): 236-48.
20. Sealy YM. Parents' food choices: Obesity among minority parents and children. *J Community Health Nurs* 2010;27(1): 1-11.
21. Sussner KM, Lindsay AC, Greaney ML, Peterson KE. The influence of immigrant status and acculturation on the development of overweight in Latino families: A qualitative study. *J Immigr Minor Health* 2008;10(6): 497-505.

22. Caprio S, Daniels SR, Drewnowski A, Kaufman FR, Palinkas LA, Rosenbloom AL, et al. Influence of race, ethnicity, and culture on childhood obesity: Implications for prevention and treatment: A consensus statement of shaping America's health and the obesity society. *Diabetes Care* 2008; *31*(11): 2211-21.
23. Berry JW. Conceptual approaches to acculturation. In Chun, KM, Organista, PB, Marin, G, editors. *Acculturation: Advances in theory, measurement, and applied research*. Washington, D.C.: American Psychological Association, 2003.
24. Grieco, EM. Race and Hispanic origin of the foreign-born population in the United States: 2007. *American Community Survey Reports 2010*. Washington, D.C: U.S. Census Bureau. Available from: <http://www.census.gov/prod/2010pubs/acs-11.pdf>
25. Allen, ML, Elliot, MN, Moarales, LS, Diamant, AL, Hambarsoomian, K., Schuster, MA. Adolescent participation in preventive health behaviors, physical activity, and nutrition: Differences across immigrant generations for Asians and Latinos compared with Whites. *Am J Public Health* 2007;*97*(2): 337-43.
26. Fuller B, Bridges M, Bein E, Jang H, Jung S, Rabe-Hesketh S, et al. The health and cognitive growth of Latino toddlers: At risk or immigrant paradox? *Matern Child Health J* 2009;*13*(6): 755-68.
27. Gordon-Larsen P, Harris KM, Ward DS, Popkin BM. Acculturation and overweight-related behaviors among Hispanic immigrants to the US: the National Longitudinal Study of Adolescent Health. *Soc Sci Med* 2003;*57*(11): 2023-34.
28. Buscemi J, Beech BM, Relyea G. Predictors of obesity in Latino children: Acculturation as a moderator of the relationship between food insecurity and body mass index percentile. *J Immigr Minor Health* 2011;*13*: 149-54.
29. Lui J, Probst JC, Harun N, Bennett KJ, Torres ME. Acculturation, physical activity, and obesity among Hispanic adolescents. *Ethn Health* 2009;*14*(5): 509-25.
30. Kallen DJ, Sussman MB, editors. *Obesity and the family: Marriage & family review*. New York: Haworth Press, 1984.

31. Schwartz MB, Puhl R. Childhood obesity: a societal problem to solve. *Obes Rev.* 2003;4:57-71.
32. Alaszewski A, Manthorpe J. Social Marxism, class conflict and health care. *Nurs Times* 1995; 91 (20): pp. 36-7.
33. DeNavas-Walt C, Proctor B, Smith J. Income, poverty, and health insurance coverage in the United States: 2010. United States Census Bureau, 2011. Available from: <http://www.census.gov/prod/2011pubs/p60-239.pdf>.
34. Evans GW, Kim P. Childhood poverty and health. Cumulative risk exposure and stress dysregulation. *Psychol Sci* 2007;18(11): 953-57.
35. Wells NM, Evans GW, Beavis A, Ong AD. Early childhood poverty, cumulative risk exposure, and body mass index trajectories through young adulthood. *Am J Public Health.* 2010 Dec;100(12): 2507-12.
36. Poulton R, Caspi A, Milne BJ, Thomson WM, Taylor A, Sears MR, et al. Association between children's experience of socioeconomic disadvantage and adult health: a life-course study. *Lancet* 2002;360(9346): 1640-5.
37. Sen B, Mennemeyer S, Gary LC. The relationship between perceptions of neighborhood characteristics and obesity among children. In Grossman M, Mocan N, editors. *Economic Aspects of Obesity*. Chicago: University of Chicago Press, 2011.
38. Reidpath DD, Burns C, Garrard J, Mahoney M, Townsend M. An ecological study of the relationship between social and environmental determinants of obesity. *Health Place* 2002;8(2): 141-45.
39. Rose, D. Economic determinants and dietary consequences of food insecurity in the United States. *J Nutr* 1999;129(2S Suppl): 517S-520S.
40. Darmon N, Drewnowski A. Does social class predict diet quality? *Am J Clin Nutr* 2008;87(5): 1107-17.

41. Balistreri, KS, Van Hook, J. Socioeconomic status and Body Mass Index among Hispanic children of immigrants and children of natives. *Am J Public Health* 2009;99(12): 2238-46.
42. Goldman N, Kimbro RT, Turra CM, Pebley AR. Socioeconomic gradients in health for white and Mexican-origin populations. *Am J Public Health* 2006;96(12): 2186-93.
43. Balistreri KS, Van Hook J. Trajectories of overweight among US school children: A focus on social and economic characteristics. *Matern Child Health J* 2011;15(5): 610-9.
44. Crawford PB, Story M, Wang MC, Ritchie LD, Sabry ZI. Ethnic issues in the epidemiology of child obesity. *Pediatr Clin North Am* 2001;48(4): 855-78.
45. Lara M, Gamboa C, Kahramanian MI, Morales LS, Bautista DE. Acculturation and Latino health in the United States: A review of the literature and its sociopolitical context. *Annu Rev Public Health*. 2005; 26:367-97.
46. Dennison BA, Tara AE, Jenkins PL. Television viewing and television in the bedroom associated with overweight risk among low-income preschool children. *Pediatr* 2002;109(6): 1028-35.
47. Elder JP, Arredondo EM, Campbell N, Baquero B, Duerksen S, Ayala G, et al. Individual, family, and community environmental correlates of obesity in Latino elementary school children. *J Sch Health* 2010;80(1): 20-30.
48. Nyberg K, Ramirez A, Gallion K. Physical activity, overweight, and obesity among Latino youth. *SaludAmerica! The Robert Wood Johnson Foundation Research Network to Prevent Obesity Among Latino Children*, 2011. Available from: <http://www.rwjf.org/files/research/73676.saludamerica.physicalactivity.pdf>
49. Harrington S. Overweight in Latino/Hispanic adolescents: Scope of the problem and nursing implications. *Pediatr Nurs* 2008;34(5): 385-96.
50. Stovitz SD, Steffen LM, Boostrom A. Participation in physical activity among normal- and overweight Hispanic and non-Hispanic white adolescents. *J Sch Health* 2008;78(1): 19-25.

51. Gallagher MR. Maternal perspectives on lifestyle habits that put children of Mexican descent at risk of obesity. *J Spec Pediatr Nurs* 2009;15(1): 16-25.
52. NCES (National Center for Education Statistics). User's manual for the ECLS-K first grade public-use data files and electronic codebook. U.S. Department of Education, Office of Educational Research and Improvement. Report number *NCES 2002-135, 2002*. Available from: <http://nces.ed.gov/pubsearch/pubsinfo.asp?pubid=2002135>
53. Kuczmarski RJ, Ogden, CL, Guo, SS, Grummer-Strawn, LM, Flegal, KM, Mei, Z, et al. CDC growth charts for the United States: Methods and development. *Vital health stat* 2002;11(246): 1-190
54. Patterson JT. *America's struggle against poverty in the twentieth century*. Cambridge, MA: Harvard University Press, 2000.
55. Cabrera NJ, Shannon JD, West J, Brooks-Gunn J. Parental interactions with Latino infants: Variation by country of origin, and English proficiency. *Child Dev* 2006;77(5): 1190-1207.
56. Liu J, Probst JC, Harun N, Bennett KJ, Torres ME. Acculturation, physical activity, and obesity among Hispanic adolescents. *Ethn Health* 2009;14(5): 509-525.
57. Amaro H, de la Torre A. Public health needs and scientific opportunities in research on Latinas. *Am J Public Health* 2002;92(4): 525-9.
58. Acevedo-Garcia D, McArdle N, Osypuk TL, Lefkowitz B, Krimgold BK. Children left behind. How metropolitan areas are failing American children. Harvard School of Public Health and Center for the Advancement of Health, 2007. Available from: [http://diversitydata-archive.org/Downloads/children\\_left\\_behind\\_final\\_report.pdf](http://diversitydata-archive.org/Downloads/children_left_behind_final_report.pdf)
59. Saelens BE, Sallis F, Black JB, Chen D. Neighborhood-based differences in physical activity: An environment scale evaluation. *Am J Public Health* 2003;93(9): 1552-8.
60. Gómez JE, Johnson BA, Selva M, Sallis JF. Violent crime and outdoor physical activity among inner-city youth. *Prev Med*. 2004;39(5): 876-81.

**Table 1**  
**Logistic regression estimates of home language, SES, TV watching, and physical activity**

	Model 1 Spring 1999		Model 2 Spring 1999		Model 3 Spring 1999		Model 4 <sup>a</sup> Spring 1999	
	OR <sup>b</sup> (95% CI)	<i>p</i> - valu e	OR <sup>b</sup> (95% CI)	<i>p</i> - value	OR <sup>b</sup> (95% CI)	<i>p</i> - value	OR <sup>b</sup> (95% CI)	<i>p</i> - value
Latino <sup>c</sup>	2.03 (1.73, 2.38)	0.00	1.24 (1.00,1.57)	0.05	1.92 (1.63, 2.27)	0.00	1.23 (0.97,1.56)	0.08
<b>Explanatory variables</b>								
SES			0.65 (0.58,0.73)	0.00			0.65 (0.58,0.74)	0.00
HL <sup>d</sup> : English & Spanish			1.52 (1.18,1.97)	0.001			1.46 (1.12,1.90)	0.005
HL: only Spanish			1.48 (1.10,2.00)	0.008			1.41 (1.0341.92)	0.026
TV watching (hours)					1.12 (1.07, 1.18)	0.00	1.09 (1.04,1.14)	0.00
Aerobic exercise					0.66 (0.57, 0.76)	0.00	0.67 (0.57,0.78)	0.00
<b>Control variables</b>								
Birth weight	1.32 (1.24, 1.41)	0.00	1.33 (1.25,1.42)	0.00	1.33 (1.25, 1.43)	0.00	1.35 (1.26,1.44)	0.00
Female	0.97 (0.84, 1.14)	0.73	0.97 (0.84,1.13)	0.73	0.98 (0.84,1.15)	0.81	0.97 (0.84,1.14)	0.76
Mother's age	1.00 (0.98, 1.01)	0.63	1.01 (1.00,1.03)	0.07	1.00 (0.99,1.02)	0.89	1.01 (1.00,1.03)	0.05
Mother's depression	1.13 (0.95, 1.35)	0.17	1.06 (0.89,1.26)	0.53	1.10 (0.91,0.31)	0.35	1.03 (0.86,1.23)	0.75
Single-parent family	1.27 (0.79, 0.91)	0.17	1.16 (0.96,1.40)	0.13	1.28 (1.05,1.56)	0.014	1.17 (0.96,1.42)	0.11
Number of siblings	0.85 (0.84, 1.14)	0.00	0.81 (0.76,0.87)	0.00	0.85 (0.79,0.91)	0.00	0.82 (0.76,0.88)	0.00

*Note:* Survey design to correct standard errors were used in all models. Unweighted sample size=8,550 children. Sample size was rounded to the nearest 10 because of restricted license requirements.

a Model 4:  $P(y_i = \text{obese}) = \text{CDF}(\beta_0 + \beta_1 \text{Latino}_i + \beta_2 \text{SES}_i + \beta_3 \text{Home language a}_i + \beta_4 \text{Home language b}_i + \beta_5 \text{TV watching}_i + \beta_6 \text{Aerobic exercise}_i + \beta_7 \text{Birth weight}_i + \beta_8 \text{female}_i + \beta_9 \text{Mother's age}_i + \beta_{10} \text{Mother's depression}_i + \beta_{11} \text{Single-parent family}_i + \beta_{11} \text{Number of sibling}_i)$ , where CDF is a Cumulative Density Function and "i" represents individual-level data.

b Coefficients reported are Odd ratio (OR), including confidence intervals (CI)

c Latino coefficient represent obesity disparities between White and Latino children

d Home language (HL)= Only English is the reference category

**Table 2 - Logistic regression estimates of home language, SES, TV watching, and physical activity**

	Model 5 <sup>a</sup> Spring 2000		Model 6 <sup>a</sup> Spring 2002		Model 7 <sup>a</sup> Spring 2004	
	OR <sup>b</sup> (95% CI)	<i>p</i> - value	OR <sup>b</sup> (95% CI)	<i>p</i> -value	OR <sup>b</sup> (95% CI)	<i>p</i> - value
Latino <sup>c</sup>	1.20 (0.96,1.50)	0.12	1.11 (0.89,1.38)	0.34	1.13 (0.90,1.42)	0.30
<b>Explanatory variables</b>						
SES	0.68 (0.60,0.76)	0.00	0.65 (0.59,0.73)	0.00	0.58 (0.51,0.66)	0.00
HL <sup>d</sup> : English & Spanish	1.73 (1.33,2.25)	0.00	1.51 (1.17,1.94)	0.002	1.43 (1.10,1.86)	0.006
HL: only Spanish	1.28 (0.92,1.79)	0.15	1.40 (1.04,1.88)	0.025	1.23 (0.90,1.67)	0.20
TV watching (hours)	1.10 (1.05,1.16)	0.00	1.10 (1.04,1.15)	0.00	1.10 (1.03,1.16)	0.002
Aerobic exercise	.69 (0.60,0.81)	0.00	0.73 (0.64,0.83)	0.00	0.79 (0.68,0.91)	0.001
<b>Control variables</b>						
Birth weight	1.27 (1.20,1.35)	0.00	1.21 (1.15,1.27)	0.00	1.16 (1.10,1.23)	0.00
Female	0.97 (0.84,1.13)	0.71	0.94 (0.82,1.07)	0.35	0.72 (0.62,0.83)	0.00
Mother's age	1.02 (1.00,1.03)	0.02	1.01 (1.00,1.03)	0.02	1.01 (0.99,1.02)	0.28
Mother's depression	0.97 (0.81,1.15)	0.66	0.97 (0.81,1.15)	0.68	1.06 (0.87,1.29)	0.54
Single-parent family	1.30 (1.08, 1.57)	0.006	1.11 (0.92,1.33)	0.28	1.15 (0.94,1.39)	0.17
Number of siblings	0.88 (0.82, 0.95)	0.02	0.87 (0.82,0.94)	0.00	0.91 (0.85,0.97)	0.006
Sample size <sup>e</sup>	7510		6700		5340	

*Note:* Survey design to correct standard errors were used in all models. The unweighted sample sizes were rounded to the nearest 10 because of restricted license requirements.

a Model 4:  $P(y_i=\text{obese}) = \text{CDF}(\beta_0 + \beta_1 \text{Latino}_i + \beta_2 \text{SES}_i + \beta_3 \text{Home language } a_i + \beta_4 \text{Home language } b_i + \beta_5 \text{TV watching}_i + \beta_6 \text{Aerobic exercise}_i + \beta_7 \text{Birth weight}_i + \beta_8 \text{female}_i + \beta_9 \text{Mother's age}_i + \beta_{10} \text{Mother's depression}_i + \beta_{11} \text{Single-parent family}_i + \beta_{11} \text{Number of sibling}_i)$ , where CDF is a Cumulative Density Function and "i" represents individual-level data

b Coefficients reported are Odd ratio (OR), including confidence intervals (CI)

c Latino coefficient represent obesity disparities between White and Latino children

d Home language (HL)= Only English is the reference category

e Sample sizes decrease over time because of ECLS-k sampling design

## Acerca de PROESA

PROESA es un centro de estudios en economía de la salud fundado por la Universidad Icesi y la Fundación Valle del Lili. Hace investigación de alta calidad y genera evidencia relevante para la orientación de las políticas públicas en protección social y economía de la salud a nivel nacional e internacional.



# PROESA

Centro de Estudios en Protección  
Social y Economía de la Salud

Universidad Icesi • Fundación Valle del Lili

Calle 18 No - 122-135, Universidad Icesi - Casa Rocha  
Teléfono: +57(2) 321-20-92  
Cali - Colombia

[www.proesa.org.co](http://www.proesa.org.co)

